

(19)



JAPANESE PATENT OFFICE

PATENT ABSTRACTS OF JAPAN

(11) Publication number: 57021320 A

(43) Date of publication of application: 04.02.1982

(51) Int. Cl. A61K 31/13
A61K 31/165

(21) Application number: 55093853
(22) Date of filing: 11.07.1980

(71) Applicant: CHUGAI PHARMACEUT CO LTD
(72) Inventor: HONDA NARIMITSU
NAGAI HIDEAKI
HINOHARA YOSHIKAZU
KOIZUMI MASUO
MURAKAMI YASUSHI
NAKANO HIDEKI


(54) BLOOD SUGAR LEVEL DEPRESSING AGENT

COPYRIGHT: (C)1982,JPO&Japio

(57) Abstract:

PURPOSE: To provide a blood sugar level depressing agent containing a specific benzamide derivative as an active component.

CONSTITUTION: An agent containing the compound of formula [R₁ and R₂ are H, alkyl, (substituted) aralkyl, or (substituted) phenyl] as an active component. The compound of formula has excellent insulin biosynthesis promoting activity and blood sugar level depressing activity. It is effective at a dose of 0.1W100mg/kg for man, and maintains the activity for ≥ 24 hr by the administration of 0.1W100mg/kg, once a day. The compound of formula can be prepared easily e.g. by reducing the corresponding m-nitrobenzoic acid amide by conventional method.



DRAFT TRANSLATION from
RISING SUN COMMUNICATIONS LTD.

(Incorporating Rotha Fullford Leopold of Canberra, Australia)

40 Bowling Green Lane, London EC1R 0NE

JAPANESE PATENT APPLICATION

No. J57-021320

A HYPOGLYCEMIC AGENT

(21) Filing no.: 55-93853

(22) Filing date: July 11, 1980.

(43) Specification published: February 4, 1982.

(72) Inventor(s): Narumitsu HONDA
c/o Chugai Pharmaceutical General Laboratories.
3-41-8, Takada, Toshima-ku, Tokyo.
Hideaki NAGAI
c/o Chugai Pharmaceutical General Laboratories.
3-41-8, Takada, Toshima-ku, Tokyo.
Masuo KOIZUMI
c/o Chugai Pharmaceutical General Laboratories.
3-41-8, Takada, Toshima-ku, Tokyo.
Yasushi MURAKAMI
c/o Chugai Pharmaceutical General Laboratories.
3-41-8, Takada, Toshima-ku, Tokyo.
Hideki NAKANO
c/o Chugai Pharmaceutical General Laboratories.
3-41-8, Takada, Toshima-ku, Tokyo.

(71) Assignee(s): Chugai Pharmaceutical KK.
5-5-1 Ukima, Kita-ku, Tokyo.

Examination request: not yet made

Number of Invention: 1

(Total 4 pages)

(51) Int.Cl. ³	Identification Code	JPO classification
A61K 31/13	ADP	6408-4C
31/165		6408-4C

Please Note- Names of Japanese firms, research laboratories and government entities, as translated are not necessarily identical with the names adopted by such organisations for international contacts. Japanese personal and surnames often permit of several readings and the ones used in this translation are not necessarily the ones preferred by their bearers. Foreign names mentioned in Japanese specifications cannot always be accurately reconstructed.


Specification

1. Title of Invention

A hypoglycemic agent.

2. Patent Claims


A hypoglycemic agent containing as effective component a compound represented by general formula



(wherein, R₁ and R₂ may be the same or different and denote a hydrogen atom, a straight-chain, branched-chain or cyclic alkyl group, an aralkyl group which can have a substituent in the nucleus, or a phenyl group which may be substituted).

3. Detailed explanation of the invention

This invention is a hypoglycemic agent containing as effective component a compound represented by general formula



(wherein, R₁ and R₂ may be the same or different and denote a hydrogen atom, a straight-chain, branched-chain or cyclic alkyl group, an aralkyl group which can have a substituent in the nucleus, or a phenyl group which may be substituted).

Among the compounds represented by aforesaid formula [I], a well known compounds are included, however, hypoglycemic action or a pharmacological action that suggests this are not described whatsoever in the prior publications describing those compounds.

The compounds represented by aforesaid formula [I] can be easily obtained for example by reduction by conventional method of corresponding meta-nitrobenzoic acid amide species as shown in the Reference Example below.

Reference Example

Into a mixed solution of 6 g isopropylamine, 15 ml triethylamine and 200 ml acetone was gradually added 18.6 g meta-nitrobenzoyl chloride under ice cooling and stirring. the mixture was stirred at the same temperature for 30 minutes and then at room temperature for one hour, thereafter, the reaction liquor was discharged into 1 litre of water, precipitated crystals were recovered by

filtration, washed with water, thereafter recrystallised, and meta-nitro-N-isoproylbenzamide (m.p. 131-132°C) 18.7 g was thereby obtained as colourless acicular crystals. Hydrogen was passed though a mixed liquor of 5.2 g of said amide, 0.5 g of 10 % palladium-carbon and 100 ml ethanol, and catalytic reduction was carried out by conventional method. After theoretical quantity hydrogen was absorbed, catalyst was eliminated, the reaction liquor was concentrated under reduced pressure, the residue was recrystallised from ethanol, and thereby meta-amino-N-isoproyl benzamide (compound 1) 4.1 g was obtained as colourless acicular crystals. m.p. 148-149°C.


Elemental analysis: as molecular formula $C_{10}H_{14}N_2O$

	C	H	N
Calculated values (%)	67.38	7.92	15.72
Measured values (%)	67.35	7.94	15.69




Compounds of Table 1 were obtained in the same way as above.











wherein, compounds 25, 27 and 29 were obtained as oily substances, the value of high mass spectra are shown in the Table and the NMR values are shown below the Table.


Table 1



[I]

Comp. No.	Substituent and position		Molecular formula	m.p. (°C)	Yield (%)	Elemental analysis value					
	R ₁	R ₂				Calc. (%)			Measured (%)		
						C	H	N	C	H	N
2	H	H	C ₇ H ₉ N ₂ O	77~78	81	61.75	5.92	20.58	61.71	5.96	20.55
3	"	CH ₃	C ₈ H ₁₀ N ₂ O	121~122	85	63.98	6.71	18.65	63.92	6.68	18.69
4	"	C ₂ H ₅	C ₉ H ₁₃ N ₂ O	70~71	76	65.83	7.37	17.06	65.72	7.28	17.19
5	"	n-C ₃ H ₇	C ₁₀ H ₁₄ N ₂ O	57~58	78	67.38	7.92	15.72	67.25	7.88	15.64
6	"	n-C ₄ H ₉	C ₁₁ H ₁₆ N ₂ O	112~113	75	68.72	8.39	14.57	68.70	8.37	14.50
7	"	sec-C ₄ H ₉	"	109~111	74		"		68.67	8.44	14.65
8	"	t-C ₄ H ₉	"	126~127	79		"		68.69	8.36	14.51
9	"	i-C ₄ H ₉	"	87~89	76		"		68.75	8.46	14.62
10	"		C ₁₃ H ₁₃ N ₂ O	147~148	84	71.52	8.31	12.83	71.58	8.35	12.76
11	"		C ₁₃ H ₁₂ N ₂ O	132~133	86	73.56	5.70	13.20	73.50	5.67	13.26
12	"		C ₁₄ H ₁₄ N ₂ O	88~89	84	74.31	6.24	12.38	74.24	6.20	13.45

Comp. No.	Substituent and position		Molecular formula	m.p. (°C)	Yield (%)	Elemental analysis value					
	R ₁	R ₂				Calc. (%)			Measured (%)		
						C	H	N	C	H	N
13	H		C ₁₆ H ₁₆ N ₂ O ₃	83~84	76	66.16	5.92	10.29	65.98	5.88	10.35
14	"		C ₁₄ H ₁₃ N ₃ O ₂	180~182	56	65.87	5.13	16.46	65.75	5.18	16.55
15	"		"	135~136	59		"		65.79	5.10	16.52
16	"		"	223~226	68		"		65.81	5.07	16.53
17	"		C ₁₃ H ₁₃ N ₃ O	151~153	79	68.70	5.77	18.49	68.64	5.79	18.43
18	"		"	130~131	71		"		68.77	5.70	18.53
19	"		"	150~151	74		"		68.75	5.67	18.42
20	"		C ₁₄ H ₁₂ N ₂ O ₂	231~233	59	65.62	4.72	10.93	65.71	4.66	11.02
21	"		C ₁₄ H ₁₄ N ₂ O	96~97	73	74.31	6.24	12.38	74.25	6.19	12.49
22	"		C ₁₅ H ₁₆ N ₂ O	94~95	80	74.97	6.71	11.66	74.92	6.75	11.61
23	"		C ₁₅ H ₁₅ N ₂ O ₂	109~110	79	70.29	6.29	10.93	70.34	6.32	10.89
24	"		C ₁₄ H ₁₃ ClN ₂ O	131~132	67	64.49	5.03	10.75	64.42	5.00	10.79

Comp. No.	Substituent and position		Molecular formula	m.p. (°C)	Yield (%)	Elemental analysis value					
	R ₁	R ₂				Calc. (%)			Measured (%)		
						C	H	N	C	H	N
25	H		C ₁₅ H ₁₆ N ₂ O	oil	82	ハイマススペクトル 240.1259			240.1246 (*1)		
26	OH ₃	OH ₃	C ₉ H ₁₂ N ₂ O	87~88	82	65.83	7.37	17.06	65.78	7.41	17.12
27	<i>m</i> -C ₃ H ₇	<i>m</i> -C ₃ H ₇	C ₁₅ H ₂₀ N ₂ O	oil	76	ハイマススペクトル 220.1571			220.1580 (*2)		
28	<i>i</i> -C ₃ H ₇	<i>i</i> -C ₃ H ₇	"	179~180	80	70.87	9.15	12.72	70.79	9.15	12.78
29	<i>m</i> -C ₄ H ₉	<i>m</i> -C ₄ H ₉	C ₁₅ H ₂₄ N ₂ O	oil	74	ハイマススペクトル 248.1883			248.1875 (*3)		
30	<i>i</i> -C ₄ H ₉	<i>i</i> -C ₄ H ₉	"	85~86	79	72.54	9.74	11.28	72.48	9.79	11.34

*1 : NMR (ODCl₃) δ : 7.55~6.40 (10H, aromatic-H, -CONH-), 3.75 (2H, s, -NH₂), 3.45 (2H, t, J=6Hz, -OH₂-), 2.75 (2H, t, J=6Hz, -CH₂-)

*2 : NMR (ODCl₃) δ : 7.35~6.50 (4H, aromatic-H), 3.90 (2H, s, -NH₂), 3.30 (4H, t, J=6Hz, (-CH₂OH₂OH₃)×2), 1.60 (4H, sextet, J=6Hz, (-CH₂CH₂OH₃)×2), 0.85 (6H, t, J=6Hz, (-OH₂OH₂CH₃)×2)

*3 : NMR (ODCl₃) δ : 7.15~6.40 (4H, aromatic-H), 4.00 (2H, s, -NH₂), 3.30 (4H, br, (-CH₂OH₂OH₂CH₃)×2), 1.40 (8H, br, (-CH₂OH₃OH₂CH₃)×2), 0.90 (6H, br, (-CH₂CH₂OH₂CH₃)×2)

The compounds of this invention obtained in this way have excellent insulin biosynthesis promotion action and hypoglycemic action, and are useful at 0.1-100 mg/kg with respect to human, and the effect thereof can be sustained for 24 hours or more by the administration of 0.1-100 mg/kg once a day.

For administration, preparations formed into desired agent form by conventional means used for normal formulation method are used.

Example 1

5-week-old DDY mice (males, body weight 25-30 g) comprising 5 animals per group were fasted for 16 hours, thereafter, aqueous solution or suspension of compounds of this invention (200 mg/kg) was orally administered, and 20 minutes later, streptozotocin 200 mg/kg was intravenously administered. Blood was collected from the heart on 24 hours later, blood sugar quantity was measured by glucose oxidase method and the plasma insulin quantity was measured by two antibody method. The measurement results are shown in Table 2.

Wherein, the compound number in the Table corresponds to the compound number of Reference Example.

Table 2

Administered compound	Blood glucose (mg/dl) mean \pm S.E.M.	Plasma Insulin (μ U/ml) mean \pm S.E.M.
Normal mouse	157 \pm 6	199 \pm 40
None (control)	386 \pm 21	43 \pm 25
1	224 \pm 19 ***	176 \pm 37 *
2	157 \pm 16 ***	153 \pm 46
3	260 \pm 33 *	213 \pm 48 *
4	248 \pm 47 *	192 \pm 54
10	263 \pm 36 *	201 \pm 38 *
12	265 \pm 32 *	253 \pm 56 *
18	166 \pm 35 ***	190 \pm 51 *
21	150 \pm 6 ***	224 \pm 30 ***
24	193 \pm 41 **	173 \pm 63
25	210 \pm 39 **	184 \pm 48 *
26	267 \pm 53	220 \pm 37 **

*: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$

Example 2

meta-aminobenzamide (compound 2)	100 pts.
calcium hydrogenphosphate	58.5 pts.
crystalline cellulose	50 pts.
corn starch	40 pts.
calcium stearate	1.5 pts.

Above components were thoroughly mixed, and tablets, 250 mg per tablet (containing 100 mg effective component) was formed by conventional method. This is used as a hypoglycemic agent.

Example 3

A 40 % aqueous solution of meta-aminobenzylbenzamide (compound 21) was prepared, and 2 ml each thereof was sealed into ampoules and sterilised. This is used as a hypoglycemic injection.

Rising Sun Communications Ltd. Terms and Conditions

Rising Sun Communications Ltd. shall not in any circumstances be liable or responsible for the accuracy or completeness of any translation unless such an undertaking has been given and authorised by Rising Sun Communications Ltd. in writing beforehand. More particularly, Rising Sun Communications Ltd. shall not in any circumstances be liable for any direct, indirect, consequential or financial loss or loss of profit resulting directly or indirectly from the use of any translation or consultation services by the customer.

Rising Sun Communications Ltd. retains the copyright to all of its' translation products unless expressly agreed in writing to the contrary. The original buyer is permitted to reproduce copies of a translation for their own corporate use at the site of purchase, however publication in written or electronic format for resale or other dissemination to a wider audience is strictly forbidden unless by prior written agreement.

⑫ 公開特許公報 (A)

昭57—21320

⑤Int. Cl.³
A 61 K 31/13
31/165

識別記号
ADP
ADP

庁内整理番号
6408—4C
6408—4C

④公開 昭和57年(1982)2月4日
発明の数 1
審査請求 未請求

(全 4 頁)

④ 血糖降下剤

①特 願 昭55—93853

②出 願 昭55(1980)7月11日

⑦発 明 者 本多成光
東京都豊島区高田3丁目41番8
号中外製薬株式会社総合研究所
内

⑧発 明 者 永井秀明
東京都豊島区高田3丁目41番8
号中外製薬株式会社総合研究所
内

⑨発 明 者 日野原好和
東京都豊島区高田3丁目41番8

号中外製薬株式会社総合研究所
内

⑩発 明 者 小泉益男
東京都豊島区高田3丁目41番8
号中外製薬株式会社総合研究所
内

⑪発 明 者 村上泰
東京都豊島区高田3丁目41番8
号中外製薬株式会社総合研究所
内

⑫出 願 人 中外製薬株式会社
東京都北区浮間5丁目5番1号

⑬代 理 人 安藤憲章

最終頁に続く


明 細 書

1. 発明の名称

血 糖 降 下 剤

2. 特許請求の範囲


一般式



(式中、 R_1 及び R_2 は同一又は異って、水素原子、直鎖・分岐鎖・環状アルキル基、核に置換基を有し得るアラルキル基又は置換基を有し得るフェニル基を示す。) で表わされる化合物を有効成分とする血糖降下剤。

3. 発明の詳細な説明

本発明は、次の一般式



[I]

(式中、 R_1 及び R_2 は同一又は異って、水素原子、直鎖・分岐鎖・環状アルキル基、核に置換基を有

し得るアラルキル基又は置換基を有し得るフェニル基を示す。) で表わされる化合物を有効成分とする血糖降下剤の発明である。

上式〔I〕で表わされる化合物の中には、公知の化合物が含まれるが、それらの記載されている先行文献には血糖降下作用ないしそれを示唆する薬理作用は全く記載されていない。

上式〔I〕で表わされる本発明の化合物は、例えば、以下の参考例に示すように、対応するメタニトロ安息香酸アミド類を常法により還元することにより容易に得ることができる。

参考例。

イソプロピルアミン6g、トリエチルアミン15ml及びアセトン200mlの混合溶液に、氷冷攪拌下、メタニトロベンゾイルクロライド18.6gを徐々に加える。同温度で30分、次いで室温で1時間攪拌後反応溶液を1ℓの水に注ぎ、析出する結晶をろ取し、水洗後再結晶して無色針状晶のメタニトロ-N-イソプロピルベンズアミド(融点131~132℃)18.7gを得た。この5.2


9、10%パラジウム-炭素0.5g及びエタノール100mlの混液に水素を通じ、常法により接触還元する。計算量の水素を吸収後触媒を除去し、反応液を減圧濃縮し、残渣をエタノールより再結晶して無色針状晶のメタアミノ-N-イソプロピルベンズアミド(化合物1)4.1gを得た。融点148~149℃。




元素分析値 分子式 $C_{10}H_{14}N_2O$ として













	C	H	N
理論値(%)	67.38	7.92	15.72
実測値(%)	67.35	7.94	15.69


上記と同様にして表1の化合物を得た。

なお、化合物25, 27及び29は油状で得られたので表中にハイマススペクトルの値を、欄外にNMRの値を記載した。

表 - 1 

化合物 No	置換基及び置換位置		分子式	融点 (℃)	収率 (%)	元 素 分 析 値					
	R ₁	R ₂				理 論 値 (%)			実 測 値 (%)		
						C	H	N	O	H	N
2	H	H	$C_7H_9N_2O$	77~78	81	61.75	5.92	20.58	61.71	5.96	20.55
3	"	CH ₃	$C_8H_{10}N_2O$	121~122	85	63.98	6.71	18.65	63.92	6.68	18.69
4	"	C ₂ H ₅	$C_9H_{12}N_2O$	70~71	76	65.83	7.37	17.06	65.72	7.28	17.19
5	"	n-C ₃ H ₇	$C_{10}H_{14}N_2O$	57~58	78	67.38	7.92	15.72	67.25	7.88	15.64
6	"	n-C ₄ H ₉	$C_{11}H_{16}N_2O$	112~113	75	68.72	8.39	14.57	68.70	8.37	14.50
7	"	sec-C ₄ H ₉	"	109~111	74	"			68.67	8.44	14.65
8	"	t-C ₄ H ₉	"	126~127	79	"			68.69	8.36	14.51
9	"	i-C ₄ H ₉	"	87~89	76	"			68.75	8.46	14.62
10	"		$C_{13}H_{18}N_2O$	147~148	84	71.52	8.31	12.83	71.58	8.35	12.76
11	"		$C_{15}H_{12}N_2O$	132~133	86	73.56	5.70	13.20	73.50	5.67	13.26
12	"		$C_{14}H_{14}N_2O$	88~89	84	74.31	6.24	12.38	74.24	6.20	13.45

No	置換基及び置換位置		分子式	融点 (°C)	収率 (%)	元素分析値					
	R ₁	R ₂				理論値(%)			実測値(%)		
						O	H	N	C	H	N
13	H		C ₁₅ H ₁₆ N ₂ O ₃	83~84	76	66.16	5.92	10.29	65.98	5.88	10.35
14	"		C ₁₄ H ₁₃ N ₃ O ₂	180~182	56	65.87	5.13	16.46	65.75	5.18	16.55
15	"		"	135~136	59	"	"	"	65.79	5.10	16.52
16	"		"	223~226	68	"	"	"	65.81	5.07	16.53
17	"		C ₁₃ H ₁₃ N ₃ O	151~153	79	68.70	5.77	18.49	68.64	5.79	18.43
18	"		"	130~131	71	"	"	"	68.77	5.70	18.53
19	"		"	150~151	74	"	"	"	68.75	5.67	18.42
20	"		C ₁₄ H ₁₂ N ₂ O ₃	231~233	59	65.62	4.72	10.93	65.71	4.66	11.02
21	"		C ₁₄ H ₁₄ N ₂ O	96~97	73	74.31	6.24	12.38	74.25	6.19	12.49
22	"		C ₁₅ H ₁₆ N ₂ O	94~95	80	74.97	6.71	11.66	74.92	6.75	11.61
23	"		C ₁₅ H ₁₆ N ₂ O ₂	109~110	79	70.29	6.29	10.93	70.34	6.32	10.89
24	"		C ₁₄ H ₁₃ O ₂ N ₂ O	131~132	67	64.49	5.03	10.75	64.42	5.00	10.79

No	置換基及び置換位置		分子式	融点 (°C)	収率 (%)	元素分析値					
	R ₁	R ₂				理論値(%)			実測値(%)		
						O	H	N	C	H	N
25	H		C ₁₅ H ₁₆ N ₂ O	oil	62	ハイマススペクトル 240.1259			(*1) 240.1246		
26	CH ₃	CH ₃	C ₉ H ₁₂ N ₂ O	87~88	82	65.83	7.37	17.06	65.78	7.41	17.12
27	n-C ₃ H ₇	n-C ₃ H ₇	C ₁₃ H ₂₀ N ₂ O	oil	76	ハイマススペクトル 220.1571			(*2) 220.1580		
28	i-C ₃ H ₇	i-C ₃ H ₇	"	179~180	80	70.87	9.15	12.72	70.79	9.15	12.78
29	n-C ₄ H ₉	n-C ₄ H ₉	C ₁₅ H ₂₄ N ₂ O	oil	74	ハイマススペクトル 248.1883			(*3) 248.1875		
30	i-C ₄ H ₉	i-C ₄ H ₉	"	85~86	79	72.54	9.74	11.28	72.48	9.79	11.34

*1: NMR (CDCl₃) δ: 7.55~6.40 (10H, aromatic-H, -CONH-), 3.75 (2H, s, -NH₂), 3.45 (2H, t, J=6Hz, -CH₂-), 2.75 (2H, t, J=6Hz, -CH₂-)

*2: NMR (ODCl₃) δ: 7.35~6.50 (4H, aromatic-H), 3.90 (2H, s, -NH₂), 3.30 (4H, t, J=6Hz, [-CH₂CH₂CH₃]₂), 1.60 (4H, sextet, J=6Hz, [-CH₂CH₂CH₃]₂), 0.85 (6H, t, J=6Hz, [-CH₂CH₂CH₃]₂)

*3: NMR (ODCl₃) δ: 7.15~6.40 (4H, aromatic-H), 4.00 (2H, s, -NH₂), 3.30 (4H, br, [-CH₂CH₂CH₂CH₃]₂), 1.40 (8H, br, [-CH₂CH₂CH₂CH₃]₂), 0.90 (6H, br, [-CH₂CH₂CH₂CH₃]₂)

このようにして得られる本発明の化合物は、優れたインスリン生合成促進作用及び血糖降下作用を有し、ヒトに対しては0.1～100mg/kgで有効で、1日1回0.1～100mg/kgの投与で24時間以上その効力を持続する。

投与に際しては、通常の製剤化に用いられる慣用手段により所望の剤形に成形された製剤が用いられる。

実施例 1.

1群5匹の5週令DDY系マウス(雄、体重25～30g)を16時間絶食後、本発明化合物(200mg/kg)の水溶液又はけん濁液を経口投与し、20分後にストレプトゾトシン200mg/kgを静脈内に投与した。24時間後に心臓から採血し、グルコースオキシダーゼ法により血中糖量を、また、二抗体法により血しょうインスリン量を測定した。測定結果を表2に例示する。

なお、表中の化合物番号は参考例の化合物番号に対応している。

投与化合物	血糖値 (mg/dl) mean ± S. E. M.	血しょうインスリン (μU/ml) mean ± S. E. M.
正常マウス	157 ± 6	199 ± 40
なし(対照)	386 ± 21	43 ± 25
1	224 ± 19***	176 ± 37*
2	157 ± 16***	153 ± 46
3	260 ± 33*	213 ± 48*
4	248 ± 47*	192 ± 54
10	263 ± 36*	201 ± 38*
12	265 ± 32*	253 ± 56*
18	166 ± 35***	190 ± 51*
21	150 ± 6***	224 ± 30**
24	193 ± 41**	173 ± 63
25	210 ± 39**	184 ± 48*
26	267 ± 53	220 ± 37**

* : P < 0.05 ** : P < 0.01 *** : P < 0.001

実施例 2.

メタアミノベンズアミド(化合物2) 100 部
リン酸水素カルシウム 58.5 部
結晶セルロース 50 部
コーンスターチ 40 部
ステアリン酸カルシウム 1.5 部

これらをよく混合し、常法により1錠250mgに打錠(有効成分100mg含有)し、血糖降下用錠剤として用いる。

実施例 3.

メタアミノ-N-ベンジルベンズアミド(化合物21)の40%水溶液を調製し、1アンプルに2mlずつ封入し、滅菌して血糖降下用注射剤として用いる。

出願人 中外製薬株式会社

代理人 安 藤 憲 章

第1頁の続き

⑦発明者 中野英樹
東京都豊島区高田3丁目41番8
号中外製薬株式会社総合研究所
内